arXiv:1412.2814v1 [math.RA] 8 Dec 2014

Color Hom-Akivis algebras, Color Hom-Leibniz algebras and Modules over Color Hom-Leibniz algebras

Ibrahima BAKAYOKO *
Département de Mathématiques,
UJNK/Centre Universitaire de N'Zérékoré,
BP : 50, N'Zérékoré, Guinea.

Momo BANGOURA†
Département de Mathématiques,
Université Gamal Abdel Nasser de Conakry,
BP : 1147, Conakry, Guinea.

Bakary MANGA‡
Département de Mathématiques et Informatique,
Université Cheikh Anta Diop de Dakar,
BP 5005 Dakar-Fann, Dakar, Senegal
and
Institut de Mathématiques et de Sciences Physiques (IMSP)
01 BP 613, Porto-Novo, Benin.

Abstract

In this paper we introduce color Hom-Akivis algebras and prove that the commutator of any color non-associative Hom-algebra structure map leads to a color Hom-akivis algebra. We give various constructions of color Hom-Akivis algebras. Next we study flexible and alternative color Hom-Akivis algebras. Likewise color Hom-Akivis algebras, we introduce non-commutative color Hom-Leibniz-Poisson algebras and presente several constructions. Moreover we give the relationship between Hom-dialgebras and Hom-Leibniz-Poisson algebras; i.e. a Hom-dialgebras give rise to a Hom-Leibniz-Poisson algebra. Finally we show that twisting a color Hom-Leibniz module structure map by a color Hom-Leibniz algebra endomorphism, we get another one.

Subject Classification: 17A30, 17D10, 16W30.
Keywords: Color Hom-Akivis algebras, Non-commutative Hom-Leibniz-Poisson algebra, Non-commutative Hom-Leibniz-Poisson modules, Hom-dialgebras.

*E-mail address: ibrahimabakayoko@yahoo.fr
†E-mail address: bangm59@yahoo.fr
‡E-mail address: bakary.manga@ucad.edu.sn, bakary.manga@imsp-uac.org

1 Introduction

Hom-algebraic structures appeared first as a generalization of Lie algebras in [3] were the authors studied q-deformations of Witt and Virasoro algebras. Other interesting Hom-type algebraic structures of many classical structures were studied as Hom-associative algebras, Hom-Lie admissible algebras and more general G-Hom-associative algebras ([23]), n-ary Hom-Nambu-Lie algebras ([7]), Hom-Lie admissible Hom-coalgebras and Hom-Hopf algebras ([22]), Hom-alternative algebras, Hom-Malcev algebras and Hom-Jordan algebras ([28]). Hom-algebraic structures were extended to the case of G-graded Lie algebras by studying Hom-Lie superalgebras, Hom-Lie admissible superalgebras in [15], color Hom-Lie algebras ([2]) and color Hom-Poisson algebras ([9]). In [31], Yuan presentes some constructions of quadratique color Hom-Lie algebras, and this is used to provide several examples. T^*-extensions and central extensions of color Hom-Lie algebras and some cohomological characterizations are established ([6]).

Akivis algebras were introduced by M. A. Akivis ([5]) as a tool in the study of some aspects of Web geometry and its connection with loop theory. These algebras were originally called "W-algebras" ([4]). Later, Hofmann and Strambach ([16]) introduce the term "Akivis algebras" for such algebraic objects.

Hom-Akivis algebras are introduced in [18], in which it is showed that the commutator of non-Hom-associative algebras lead to Hom-Akivis algebras. It is also proved that Hom-Akivis algebras can be obtain from Akivis algebras by twisting along algebra endomorphisms, and that the class of Hom-Akivis algebras is closed under self-morphisms. The connection between Hom-Akivis algebras and Hom-alternative algebras is given.

A non-commutative version of Lie algebras were introduced by Loday in [21]. The bracket of the so-called Leibniz algebras satisfies the Leibniz identity; which, combined with skew-symmetry, is a variation of the Jacobi identity. hence Lie algebras are skew-symmetric Leibniz algebras. In the Leibniz setting, the objects that play the role of associative algebras are called dialgebras, which were introduced by Loday in [20]. Dialgebras have two associative binary operations that satisfy three additional associative-type axioms. The relationship between dialgebras and Leibniz algebras are analyzed in [13]. It is proved in [26] that from any Leibniz algebra L one can construct a Leibniz-Poisson algebra A and the properties of L are close to the properties of A. Leibniz algebras were extended to Hom-Leibniz algebras in [24]. A (co)homology theory of Hom-Leibniz algebras and an initial study of universal central extensions of Hom-Leibniz algebras was given in [14]. The representation of Hom-Leibniz algebras are introduced in [12] and the homology of Hom-Leibniz algebras are computed. Recently Leibniz superalgebras were studied in [1]. The Hom-dialgebras are introduced in [30] as an extention of some work of Loday ([20]). In a Hom-dialgebra, there are two binary operations that satisfy five α-twisted associative-type axioms. Each Hom-Lie algebra can be thought of as a Hom-Leibniz algebra. Likewise, to every Hom-associative algebra is associated a Hom-dialgebra in which both binary operations are equal to the original one.

The purpose of this paper is to study color Hom-Akivis algebras (graded version of Hom-Akivis algebras ([18])), color non-commutative-Hom-Leibniz-Poisson algebras and modules over color Hom-Leibniz algebras.

Section 2 is dedicated to the background material on graded vector spaces, bicharac-

ter and non-Hom-associative algebras. In section 3, we define color Hom-Akivis algebras which is the graded version of Hom-Akivis algebras. Various properties of color Hom-Akivis algebras are studied. More precisely, we show that the commutator of color non-Hom-associative algebra structure map lead to color Hom-Akivis algebras (Theorem 3.1). Next, we provide some constructions of color Hom-Akivis algebras; on the one hand from color Akivis algebras (Proposition 3.1) and on the other hand from a given color Hom-Akivis algebras (Theorem 3.2). Flexible color Hom-Akivis algebras and Hom-alternative color Hom-Akivis algebras are also exposed (Theorem 3.3). Section 4 being devoted to the study of Color Hom-Leibniz algebras, we define in it color Hom-Leibniz algebras and give basic properties (Proposition 4.1). Next, we introduce non-commutative Hom-Leibniz-Poisson algebras (in brief color NHLP-algebras). As in the previous section on color Hom-Akivis algebras, we give several twisting of non-commutative Hom-Leibniz-Poisson algebras (Theorem 4.1). Finally we give the relationship between Hom-dialgebras and Hom-Leibniz-Poisson algebras (Theorem 4.3). In section 5, we prove that the Yau's twisting of module Hom-algebras ([29, Lemma 2.5]) works for modules over color Hom-Leibniz algebras (Theorem 5.1) likewise it runs for modules over Hom-Lie algebras ([10]), modules over color Hom-Lie algebras ([9]) and modules over color Hom-Poisson algebras ([9]).

Throughout this paper, $\mathbf{K}$ denotes a field of characteristic 0 and G is an abelian group.

2 Preliminaries

Definition 2.1. *A vector space V is said to be a G-graded if, there exist a family $(V_a)_{a\in G}$ of vector subspaces of V such that $V = \bigoplus_{a\in G} V_a$. An element $x \in V$ is said to be homogeneous of degree $a \in G$ if $x \in V_a$. We denote $\mathcal{H}(V)$ the set of all homogeneous elements in V.*

Definition 2.2. *Let $V = \bigoplus_{a\in G} V_a$ and $V' = \bigoplus_{a\in G} V'_a$ be two G-graded vector spaces, and $b \in G$. A linear map $f : V \to V'$ is said to be homogeneous of degree b if $f(V_a) \subseteq V'_{a+b}$, for all $a \in G$. If, f is homogeneous of degree zero i.e. $f(V_a) \subseteq V'_a$ holds for any $a \in G$, then f is said to be even.*

Definition 2.3. *An algebra (A, μ) is said to be G-graded if its underlying vector space is G-graded and if, furthermore, $\mu(A_a, A_b) \subseteq A_{a+b}$, for all $a, b \in G$. Let A' be another G-graded algebra. A morphism $f : A \to A'$ of G-graded algebras is by definition an algebra morphism from A to A' which is, in addition an even mapping.*

Definition 2.4. *Let G be an abelian group. A map $\varepsilon : G \times G \to \mathbf{K}^*$ is called a skew-symmetric bicharacter on G if the following identities hold: for all $a, b, c \in G$,*

(i) $\varepsilon(a, b)\varepsilon(b, a) = 1$;

(ii) $\varepsilon(a, b + c) = \varepsilon(a, b)\varepsilon(a, c)$;

(iii) $\varepsilon(a + b, c) = \varepsilon(a, c)\varepsilon(b, c)$.

Definition 2.5. *1. A multiplicative color non-Hom-associative (i.e. non necessarily Hom-associative) algebra is a quadruple $(A, \mu, \varepsilon, \alpha_A)$ such that*

(a) *A is G-graded vector space;*

(b) *$\mu : A \otimes A \to A$ is an even bilinear map;*

(c) *$\varepsilon : G \otimes G \to K^*$ is a bicharacter;*

(d) *α_A is an endomorphism of (A, μ) (multiplicativity).*

2. *(A, μ, α) is said to be ε-skew-symmetric (resp. ε-commutative) if, for any $x, y \in A$, $\mu(x, y) = -\varepsilon(x, y)\mu(y, x)$ (resp. $\mu(x, y) = \varepsilon(x, y)\mu(y, x)$).*

3. *Let (A, μ, α_A) be color non-Hom-associative algebra. For any $x, y, z \in A$, the Hom-associator is defined by*

$$as(x, y, z) = \mu(\mu(x, y), \alpha_A(z)) - \mu(\alpha_A, \mu(y, z)). \tag{1}$$

Then $(A, \mu, \varepsilon, \alpha_A)$ is said to be

(a) *Hom-flexible, if $as(x, y, x) = 0$;*

(b) *Hom-alternative, if $as(x, y, z)$ is ε-skew-symmetric in x, y, z.*

3 Color Hom-akivis algebras

Hom-Akivis algebras was introduced in [18] as a twisted generalization of Akivis algebras. In the following, we generalize some results. We prove that the commutator of a color non-Hom-associative algebra leads to a color Hom-Akivis algebra. Some properties of color Hom-Akivis algebras are studied.

Definition 3.1. *A color Hom-Akivis algebra is quintuple $(A, [-,-], [-,-,-], \varepsilon, \alpha_A)$ consisting of a G-graded vector space A, an even ε-skew-symmetric bilinear map $[-,-]$, an even trilinear map $[-,-,-]$ and an even endomorphism $\alpha_A : A \to A$ such that for all $x, y, z \in \mathcal{H}(A)$,*

$$\oint \varepsilon(z, x)[[x, y], \alpha_A(z)] = \oint \varepsilon(z, x)\Big([x, y, z] - \varepsilon(x, y)[y, x, z]\Big). \tag{2}$$

If in addition, α_A is an endomorphism with respect to $[-,-]$ and $[-,-,-]$, the color Hom-Akivis algebra A is said to be multiplicative.

Theorem 3.1. *Let $(A, \mu, \varepsilon, \alpha_A)$ be a color non-Hom-associative algebra. Define the maps $[-,-] : A \otimes A \to A$ and $[-,-,-]_{\alpha_A} : A \otimes A \otimes A \to A$ as follows: for all $x, y, z \in A$,*

$$\begin{aligned} [x, y] &:= \mu(x, y) - \varepsilon(x, y)\mu(y, x), \\ [x, y, z]_{\alpha_A} &:= as(x, y, z). \end{aligned}$$

Then $(A, [-,-], [-,-,-]_{\alpha_A}, \varepsilon, \alpha_A)$ is a multiplicative color Hom-Akivis algebra.

Proof. Color Hom-Akivis identity (2) is proved by direct computation. For any homogeneous elements $x, y, z \in A$, we have

$$\begin{aligned} \varepsilon(z, x)[[x, y], \alpha_A(z)] &= \varepsilon(z, x)(xy)\alpha_A(z) - \varepsilon(y, z)\alpha_A(z)(xy) \\ &\quad -\varepsilon(z, x)\varepsilon(x, y)(yx)\alpha_A(z) + \varepsilon(x, y)\varepsilon(y, z)\alpha_A(z)(yx). \end{aligned}$$

And,

$$\begin{aligned}\varepsilon(z,x)([x,y,z]-\varepsilon(x,y)[y,x,z]) &= \varepsilon(z,x)(xy)\alpha_A(z)-\varepsilon(z,x)\alpha_A(x)(yz)\\ &\quad -\varepsilon(z,x)\varepsilon(x,y)(yx)\alpha_A(z)+\varepsilon(z,x)\varepsilon(x,y)\alpha_A(y)(xz).\end{aligned}$$

Expanding $\oint \varepsilon(z,x)[[x,y],\alpha_A(z)]$ and $\oint \varepsilon(z,x)([x,y,z]-\varepsilon(x,y)[y,x,z])$, one gets (2) and so the conclusion holds. The multiplicativity follows from that of $(A,\mu,\varepsilon,\alpha_A)$. □

Definition 3.2. *Let $(A,[-,-],[-,-,-],\varepsilon,\alpha_A)$ and $(\tilde{A},\{-,-\},\{-,-,-\},\varepsilon,\alpha_{\tilde{A}})$ be two color Hom-Akivis algebras. A morphism $f: A\to\tilde{A}$ of color Hom-Akivis algebras is an even linear map of G-graded vector spaces A and $\tilde{A}$ such that*

$$\begin{aligned}f([x,y]) &= \{f(x),f(y)\},\\ f([x,y,z]) &= \{f(x),f(y),f(z)\}.\end{aligned}$$

For example, if we take $(A,[-,-],[-,-,-],\varepsilon,\alpha_A)$ as a multiplicative color Hom-Akivis algebra, then the twisting self-map α_A is itself an endomorphism of $(A,[-,-],[-,-,-],\varepsilon)$.

The following theorem is the color version of Theorem 4.4 in [18].

Theorem 3.2. *Let $A_{\alpha_A}=(A,[-,-],[-,-,-],\varepsilon,\alpha_A)$ be a color Hom-Akivis algebra and $\beta: A\to A$ an even endomorphism. Then, for any integer $n\geq 1$, $A_{\beta^n}=(A,[-,-]_{\beta^n},[-,-,-]_{\beta^n},\varepsilon,\beta^n\circ\alpha_A)$ is a color Hom-Akivis algebra, where for all $x,y,z\in\mathcal{H}(A)$,*

$$\begin{aligned}[x,y]_{\beta^n} &:= \beta^n([x,y]),\\ [x,y,z]_{\beta^n} &:= \beta^{2n}([x,y,z]).\end{aligned}$$

Moreover, if A_{α_A} is multiplicative and β commutes with α_A, then A_{β^n} is multiplicative.

Proof. It is clear that $[-,-]_{\beta^n}$ and $[-,-,-]_{\beta^n}$ are bilinear and trilinear respectively. It is also clear that the ε-skew-symmetry of $[-,-]_{\beta^n}$ comes from that of $[-,-]$. It remains to prove the color Hom-Akivis identity (2) and the multiplicativity of A_{β^n}. For any $x,y,z\in\mathcal{H}(A)$, we have,

$$\begin{aligned}\oint \varepsilon(z,x)[[x,y]_{\beta^n},(\beta^n\circ\alpha_A)(z)]_{\beta^n} &= \oint \varepsilon(z,x)\beta^n[\beta^n[x,y],(\beta^n\circ\alpha_A)(z)]\\ &= \beta^{2n}\oint \varepsilon(z,x)([x,y,z]-\varepsilon(x,y)[y,x,z])\\ &= \oint \varepsilon(z,x)(\beta^{2n}[x,y,z]-\varepsilon(x,y)\beta^{2n}[y,x,z])\\ &= \oint \varepsilon(z,x)([x,y,z]_{\beta^{2n}}-\varepsilon(x,y)[y,x,z]_{\beta^{2n}}).\end{aligned}$$

The multiplicativity is proved similarly as in [18, Theorem 4.4]. □

Corollary 3.1. *Let $(A,[-,-],[-,-,-],\varepsilon)$ be a color Akivis algebra and $\beta: A\to A$ an even endomorphism. Then $(A,[-,-]_{\beta^n},[-,-,-]_{\beta^n},\varepsilon,\alpha_A)$ is a multiplicative color Hom-Akivis algebra.*

Moreover, suppose that $(\tilde{A},\{-,-\},\{-,-,-\},\varepsilon)$ is another color Akivis algebra and $\tilde{\alpha}_A$ an even endomorphism of $\tilde{A}$. If $f: A\to\tilde{A}$ is a morphism of color Akivis algebras such that $f\circ\alpha_A=\alpha_{\tilde{A}}\circ f$, then $f:(A,[-,-],[-,-,-],\varepsilon,\alpha_A)\to(\tilde{A},\{-,-\},\{-,-,-\},\varepsilon,\alpha_{\tilde{A}})$ is a morphism of multiplicative color Hom-Akivis algebras.

Proof. It is similar to that of [18, Corollary 4.5]. □

The above construction of color Hom-Akivis algebras is used in [18], in the case of Hom-Akivis algebras, to produce examples of Hom-Akivis algebras.

Proposition 3.1. *Let* $(A, [-,-], [-,-,-], \varepsilon)$ *be a color Akivis algebra and* $\beta : A \to A$ *an even endomorphism. Define* $[-,-]_{\beta^n}$ *and* $[-,-,-]_{\beta^n}$ *by: for all* $x, y, z \in \mathcal{H}(A)$,

$$\begin{aligned} [x,y]_{\beta^n} &:= \beta([x,y]_{\beta^{n-1}}), \\ [x,y,z]_{\beta^n} &:= \beta^2([x,y,z]_{\beta^{n-1}}), \end{aligned}$$

Then $(A, [-,-]_{\beta^n}, [-,-,-]_{\beta^n}, \varepsilon, \alpha_A)$ *is a multiplicative color Hom-Akivis algebra.*

Proof. It is proved recurrently by applying Corollary 3.1. The reader may also see [18, Theorem 4.8] for the proof. □

Definition 3.3. *A color Hom-Akivis algebra* $(A, [-,-], [-,-,-], \varepsilon), \alpha_A$ *is said to be*

1. *Hom-flexible, if* $[x,y,x] = 0$ *for all* $x, y \in A$ *;*
2. *Hom-alternative, if* $[-,-,-]$ *is* ε*-alternating (i.e.* $[-,-,-]$ *is* ε*-skew- symmetric for any pair of variables).*

Theorem 3.3. *Let* $(A, \mu, \varepsilon, \alpha_A)$ *be a color non-Hom-associative algebra and the quintuple* $(A, [-,-], as(-,-,-), \varepsilon, \alpha_A)$ *its associated color Hom-Akivis algebra (as in Theorem 3.1). Then we have the following:*

1. *if* $(A, \mu, \varepsilon, \alpha_A)$ *is Hom-flexible, then* $(A, [-,-], as(-,-,-), \varepsilon, \alpha_A)$ *is Hom-flexible;*
2. *if* $(A, \mu, \varepsilon, \alpha_A)$ *is Hom-alternative, then so is* $(A, [-,-], as(-,-,-), \varepsilon, \alpha_A)$.

Proposition 3.2. *Let* $\mathcal{A}_{\alpha_A} = (A, [-,-], [-,-,-], \varepsilon, \alpha_A)$ *be a Hom-flexible color Hom-Akivis algebra. Then we have*

$$\oint \varepsilon(z,x) = \oint \Big(\varepsilon(z,x) + \varepsilon(x,y)\varepsilon(y,z)\Big)[x,y,z]. \tag{3}$$

In particular, $\mathcal{A}_{\alpha_A}$ *is a color Hom-Lie algebra if and only if*

$$\oint \Big(\varepsilon(z,x) + \varepsilon(x,y)\varepsilon(y,z)\Big)[x,y,z] = 0. \tag{4}$$

Proof. By expanding the left hand side of color Hom-Akivis identity (2), and using the Hom-flexibility we get (3). □

For more details on color Hom-Lie algebras, see [2], [9], [31].

4 Color Hom-Leibniz algebras

In this section, we write NHLP-algebra (resp. NLP-algebra) for non-commutative Hom-Leibniz-Poisson algebra (resp. non-commutative Leibniz-Poisson algebra). Interrelation between Hom-dialgebras and NHLP-algebras are presented.

4.1 Color NHLP-algebras

This subsection is devoted to color Hom-Leibniz algebras. Color NHLP-algebras are introduced and their various twisting are given as well as examples.

Definition 4.1. *A color Hom-Leibniz algebra is a quadruple $(L, \mu, \varepsilon, \alpha_L)$, consisting of a graded vector space L, an even bilinear map $\mu : L \otimes L \to L$, a bicharacter $\varepsilon : G \otimes G \to \mathbf{K}^*$ and an even linear space homomorphism $\alpha_L : L \to L$ satisfying*

$$\mu\Big(\alpha_L(x), \mu(y,z)\Big) = \mu\Big(\mu(x,z), \alpha(y)\Big) + \varepsilon(x,y)\mu\Big(\alpha(x), \mu(y,z)\Big). \tag{5}$$

A morphism of color Hom-Leibniz algebras is an even linear map which preserves the structures.

Remark 4.1. *1. When G is an abelian group with trivial grading, we obtain the following Hom-Leibniz identity ([18]):*

$$\mu\Big(\alpha_L(x), \mu(y,z)\Big) = \mu\Big(\mu(x,y), \alpha(z)\Big) + \mu\Big(\alpha(y), \mu(x,z)\Big). \tag{6}$$

2. The original definition of Hom-Leibniz algebras ([24]) is related to the identity

$$\mu\Big(\mu(x,y), \alpha_L(z)\Big) = \mu\Big(\alpha(x), \mu(y,z)\Big) + \mu\Big(\mu(x,z), \alpha(y)\Big) \tag{7}$$

which is expressed in terms of (right) adjoint homomorphisms $ad_x y : \mu(x,y)$ of (A, μ, α_A). This justifies the terms of "(right) Hom-Leibniz algebra" that could be used for the Hom-Leibniz algebras defined in [24]. The dual of (6) is (7) and in this paper we consider only left color Hom-Leibniz algebras.

We have the following result.

Proposition 4.1. *Let $(L, \mu, \varepsilon, \alpha_L)$ be a color Hom-Leibniz algebra. Then*

$$\Big(x \cdot y + \varepsilon(x,y) y \cdot x\Big)\alpha_L(z) = 0, \tag{8}$$

$$[x \cdot y, \alpha_L(z)] + \varepsilon(x,y)[\alpha_L(y), x \cdot z] = \alpha_L(x) \cdot [y,z], \tag{9}$$

where we have putted $\mu(x,y) = x \cdot y$ and $[x,y] = x \cdot y - \varepsilon(x,y) y \cdot x$.

Proof. The identity (6) implies that

$$(x \cdot y) \cdot \alpha_L(z) = \alpha_L(x) \cdot (y \cdot z) - \varepsilon(x,y)\alpha_L(y) \cdot (x \cdot z)$$

Likewise, interchanging x and y, we have

$$(y \cdot x) \cdot \alpha_L(z) = \alpha_L(y) \cdot (x \cdot z) - \varepsilon(y,x)\alpha_L(x) \cdot (y \cdot z)$$

Then, multiplying the second equality by $\varepsilon(x,y)$, and adding memberwise with the first one, we obtain (8). Next, by direct computation, we have

$$[x \cdot y, \alpha_L(z)] + \varepsilon(x,y)[\alpha_L(y), x \cdot z] = (x \cdot y) \cdot \alpha_L(z) - \varepsilon(x,z)\varepsilon(y,z)\alpha_L(z) \cdot (x \cdot y)$$

$$
\begin{aligned}
&+\varepsilon(x,y)\alpha_L(y)\cdot(x\cdot z)-\varepsilon(y,z)(x\cdot z)\cdot\alpha_L(y)\\
&=\alpha_L(x)\cdot(y\cdot z)-\varepsilon(x,z)\varepsilon(y,z)((z\cdot x)\cdot\alpha_L(y)\\
&\quad+\varepsilon(z,x)\alpha_L(x)\cdot(z\cdot y))-\varepsilon(y,z)(x\cdot z)\cdot\alpha_L(y) \text{ (by (5))}\\
&=\alpha_L(x)\cdot(y\cdot z)-\varepsilon(y,z)\alpha_L(x)\cdot(z\cdot y)\quad\text{(by (8))}\\
&=\alpha_L(x)\cdot[y,z]. \qquad \square
\end{aligned}
$$

Now we define color NHLP-algebras which is the graded and Hom-version of NLP-algebras ([13]).

Definition 4.2. *A color NHLP-algebra is a G-graded vector space P together with two even bilinear maps $[-,-]:P\otimes P\to P$ and $\mu:P\otimes P\to P$, a bicharacter $\varepsilon:G\otimes G\to\mathbf{K}^*$ and $\alpha_P:P\to P$ an even linear map such that, for any $x,y,z\in P$,*

i) $(P,[\cdot,\cdot],\varepsilon,\alpha_P)$ *is a color Hom-Leibniz-Poisson algebra i.e.*

$$[\alpha_P(x),[y,z]]=[[x,y],\alpha_P(z)]+\varepsilon(x,y)[\alpha_P(y),[x,z]], \tag{10}$$

ii) $(P,\mu,\varepsilon,\alpha_P)$ *is a color Hom-associative algebra i.e.*

$$\mu\big(\alpha(x),\mu(y,z)\big)=\mu\big(\mu(x,y),\alpha(z)\big)\ \textit{(Hom-associativity)}\ ; \tag{11}$$

iii) and the following identity holds:

$$[\alpha_P(x),\mu(y,z)]=\mu([x,y],\alpha_P(z))+\varepsilon(x,y)\mu(\alpha_P(y),[x,z]). \tag{12}$$

If in addition, α_P is an endomorphism with respect to μ and $[-,-]$, we say that $(P,\mu,[\cdot,\cdot],\varepsilon,\alpha_P)$ is a multiplicative color NHLP-algebra.

Remark 4.2. *When the Hom-associative product μ is ε-commutative, then $(A,\mu,[-,-]),\varepsilon,\alpha_A)$ is said to be a commutative color Hom-Leibniz Poisson algebra.*

Example 4.1. *1. Any color Hom-Poisson algebra is a color NHLP-algebra algebra.*

2. Any color Hom-Leibniz algebra is a color NHLP-algebra.

3. If P is a Leibniz-Poisson algebras (viewed as Hom-Leibniz-Poisson algebras with trivial twisting and trivial grading), then the vector space $P\otimes P$ is a NHLP-algebra with the operations

$$(x_1\otimes x_2)(y_1\otimes y_2)=x_1y_1\otimes y_1y_2,$$

$$[x_1\otimes x_2,y_1\otimes y_2]=[[x_1,x_2],y_1]\otimes y_2+y_1\otimes[[x_1,x_2],y_2].$$

The proposition below is a direct consequence of the Definition 4.2.

Proposition 4.2. *Let $(P,[\cdot,\cdot],\mu,\varepsilon,\alpha)$ be a color NHLP-algebra. Then $(P,[\cdot,\cdot],\mu^{op},\varepsilon,\alpha)$ and $(P,k\mu,k[-,-],\alpha_P)$ are also color NHLP-algebras, with $\mu^{op}(x,y)=\mu(y,x)$ and $k\in\mathbf{K}^*$.*

It is well known in Hom-algebras setting that one can obtain Hom-algebra structures from an ordinary one and an endomorphism. The following theorem, which gives a way to construct a color NHLP-algebras from color NLP-algebras and an endomorphism, is a similar result.

Theorem 4.1. *Let $(A, \mu, [-,-], \varepsilon)$ be a color NLP-algebra and $\alpha_A : A \to A$ an even endomorphism. Then, for any integer $n \geq 1$, $(A, \mu_{\alpha_A^n}, [-,-]_{\alpha_A^n}, \varepsilon, \alpha_A^n)$ is a multiplicative color NHLP-algebra, where for all $x, y, z \in \mathcal{H}(A)$,*

$$\begin{aligned} \mu_{\alpha_A^n}(x,y) &:= \alpha_A^n(\mu(x,y)), \\ [x,y]_{\alpha_A^n}(x,y) &:= \alpha_A^n([x,y]). \end{aligned}$$

Moreover, suppose that $(\tilde{A}, \tilde{\mu}, \{-,-\}, \varepsilon)$ is another color NLP-algebra and $\tilde{\alpha}_A$ an even endomorphism of $\tilde{A}$. If $f : A \to \tilde{A}$ is a morphism of color NLP-algebras such that $f \circ \alpha_A = \alpha_{\tilde{A}} \circ f$, then $f : (A, \mu, [-,-], \varepsilon, \alpha_A) \to (\tilde{A}, \tilde{\mu}, \{-,-,-\}, \varepsilon, \alpha_{\tilde{A}})$ is a morphism of multiplicative color NHLP-algebras.

Proof. We need to show that $(A, \mu_{\alpha_A^n}, [-,-]_{\alpha_A^n}, \varepsilon, \alpha_A^n)$ satisfies relations (10) and (12). We have, for any $x, y, z \in A$,

$$\begin{aligned} [\alpha_A^n(x), \mu_{\alpha_A^n}(y,z)]_{\alpha_A^n} &= \alpha_A^n([\alpha_A^n(x), \alpha_A^n\mu(y,z)]) \\ &= \alpha_A^{2n}([x, \mu(y,z)]) \\ &= \alpha_A^{2n}([[x,y],z] + \varepsilon(x,y)[y,[x,z]]) \\ &= \alpha_A^n([\alpha_A^n([x,y]), \alpha_A^n(z)] + \varepsilon(x,y)[\alpha_A^n(y), \alpha_A^n([x,z])]) \\ &= [[x,y]_{\alpha_A^n}, \alpha_A^n(z)]_{\alpha_A^n} + \varepsilon(x,y)[\alpha_A^n(y), [x,z]]_{\alpha_A^n}. \end{aligned}$$

The compatibility condition (12) is pointed out similarly and the second assertion is proved as in the case of Theorem 3.2. □

Example 4.2. *The commutative Leibniz-Poisson algebra used in this example is given in [27], [26]. Let $(A, [-,-])$ is a Leibniz algebra over a commutative infinite field $\mathbf{K}$ and let*

$$\tilde{A} = A \oplus \mathbf{K}$$

be a vector space with multiplications $\cdot$ and $\{-,-\}$ defined as: for $x, y \in A$ and $a, b \in \mathbf{K}$,

$$(x+a)\cdot(y+b) = (bx+ay) + ab \quad \text{and} \quad \{x+a, y+b\} = [x,y].$$

Consider the homomorphism $\alpha_{\tilde{A}} = \alpha_A \oplus Id_{\mathbf{K}} : \tilde{A} \to \tilde{A}$, then $(\tilde{A}, \cdot, \{-,-\}, \alpha_{\tilde{A}})$ is a commutative color Hom-Leibniz-Poisson algebra (with trivial grading).

The next result gives a procedure to produce color NHLP-algebras from given one.

Theorem 4.2. *Let $(P, \mu, [-,-], \varepsilon, \alpha_P)$ be a color NHLP-algebra and $\beta : P \to P$ an even endomorphism. Then $(P, \mu_{\beta^n}, [-,-]_{\beta^n}, \varepsilon, \beta^n \circ \alpha_P)$ is a color NHLP-algebra, with*

$$\begin{aligned} \mu_{\beta^n}(x,y) &:= \beta^n(\mu(x,y)), \\ [x,y]_{\beta^n}(x,y) &:= \beta^n([x,y]), \end{aligned}$$

for all $x, y, z \in \mathcal{H}(P)$.

Proof. We need to show that $(P, \mu_{\beta^n}, [-,-]_{\beta^n}, \varepsilon, \beta^n \circ \alpha_P)$ satisfies relations (10) and (12). We have, for any $x, y, z \in \mathcal{H}(P)$,

$$\begin{aligned}
[(\beta^n \circ \alpha_P)(x), [y,z]_{\beta^n}]_{\beta^n} &= \beta^n[(\beta^n \circ \alpha_P)(x), \beta^n[y,z]] \\
&= \beta^{2n}([\alpha_P(x), [y,z]]) \\
&= \beta^{2n}([[x,y], \alpha_P(z)] + \varepsilon(x,y)[\alpha_P(y), [x,z]]) \\
&= \beta^n[\beta^n[x,y], (\beta^n \circ \alpha_P)(z)] + \varepsilon(x,y)[(\beta^n \circ \alpha_P)(y), \beta^n[x,z]] \\
&= [[x,y]_{\beta^n}, (\beta^n \circ \alpha_P)(z)]_{\beta^n} + \varepsilon(x,y)[(\beta^n \circ \alpha_P)(y), [x,z]_{\beta^n}]_{\beta^n}.
\end{aligned}$$

Next,

$$\begin{aligned}
[(\beta^n \circ \alpha_P)(x), \mu_{\beta^n}(y,z)]_{\beta^n} &= \beta^n[(\beta^n \circ \alpha_P)(x), \beta^n\mu(y,z)] \\
&= \beta^{2n}[\alpha_P(x), \mu(y,z)] \\
&= \beta^{2n}([[x,y], \alpha_P(z)] + \varepsilon(x,y)[\alpha_P(y), [x,z]]) \\
&= \beta^n([\beta^n[x,y], (\beta^n \circ \alpha_P)(z)] + \varepsilon(x,y)[(\beta^n \circ \alpha_P)(y), \beta^n[x,z]]) \\
&= [[x,y]_{\beta^n}, (\beta^n \circ \alpha_P)(z)]_{\beta^n} + \varepsilon(x,y)[(\beta^n \circ \alpha_P)(y), [x,z]_{\beta^n}]_{\beta^n}. \quad \square
\end{aligned}$$

Proposition 4.3. *Let $(P, \mu, [-,-], \varepsilon)$ be a color NLP-algebra and $\beta : P \to P$ an even endomorphism. Define μ_{β^n} and $[-,-]_{\beta^n}$ by*

$$\begin{aligned}
\mu_{\beta^n}(x,y) &:= \beta(\mu_{\beta^{n-1}}(x,y)), \\
[x,y]_{\beta^n} &:= \beta^n([x,y]_{\beta^{n-1}}),
\end{aligned}$$

for all $x, y, z \in \mathcal{H}(P)$. Then $(P, \mu_{\beta^n}, [-,-]_{\beta^n}, \varepsilon, \alpha_P)$ is a multiplicative color NHLP-algebra.

4.2 NHLP-algebras and Dialgebras

It is well known that Hom-dialgebras give rise to Hom-Leibniz algebra ([30]). In this subsection, we give a similar result for NHLP-algebras.

Definition 4.3 ([30])**.** *A Hom-dialgebra is a quadruple $(D, \dashv, \vdash, \alpha_D)$, where D is a* **K***-vector space, $\dashv, \vdash : D \otimes D \to D$ are bilinear maps and $\alpha_D : D \to D$ a linear map such that the following five axioms are satisfied for $x, y, z \in D$:*

$$\begin{aligned}
(x \vdash y) \dashv \alpha_D(z) &= \alpha_D(x) \vdash (y \dashv z), \\
\alpha_D(x) \dashv (y \dashv z) &= (x \dashv y) \dashv \alpha_D(z) = \alpha_D(x) \dashv (y \vdash z) \\
(x \dashv y) \vdash \alpha_D(z) &= = \alpha_D(x) \vdash (y \vdash z) = (x \vdash y) \vdash \alpha_D(z)
\end{aligned}$$

We say that a Hom-dialgebra $(D, \dashv, \vdash, \alpha_D)$ is Hom-associative if it so for the operations $\vdash$ and $\dashv$. So any Hom-dialgebra is Hom-associative.

The following result connects Hom-dialgebras and Hom-Leibniz-Poisson algebras.

Theorem 4.3. *Let $(D, \dashv, \vdash, \alpha_D)$ be a Hom-dialgebra. Define the bilinear map $[-,-] : D \otimes D \to D$ by setting*

$$[x,y] := x \vdash y - y \dashv x. \tag{13}$$

Then $(D, \dashv, [-,-], \alpha_D)$ is a Hom-Leibniz-Poisson algebra.

Proof. We write down all twelve terms involved in the Hom-Leibniz identity (6) :

$$\begin{aligned}
[\alpha_D(x),[y,z]] &= \alpha_D(x)\vdash(y\vdash z)-(y\vdash z)\dashv\alpha_D(x)-\alpha_D(x)\vdash(z\dashv y)+(z\dashv y)\dashv\alpha_D(x)\\
[\alpha_D(y),[x,z]] &= \alpha_D(y)\vdash(x\vdash z)-(x\vdash z)\dashv\alpha_D(y)-\alpha_D(y)\vdash(z\dashv x)+(z\dashv x)\dashv\alpha_D(y)\\
[[x,y],\alpha_D(z)] &= (x\vdash y)\vdash\alpha_D(z)-\alpha_D(z)\dashv(x\vdash y)-(y\dashv x)\vdash\alpha_D(z)+\alpha_D(z)\dashv(y\dashv x).
\end{aligned}$$

Using the Hom-dialgebra axioms in Definition 4.3, it is readily seen that (6) holds. Next,

$$\begin{aligned}
[\alpha_D(x),y\dashv z]-[x,y]\dashv\alpha_D(z)-\alpha_D(y)\dashv[x,z] = {}&\alpha_D(x)\vdash(y\dashv z)-(y\dashv z)\dashv\alpha_D(x)\\
&-(x\vdash y)\dashv\alpha_D(z)+(y\dashv x)\dashv\alpha_D(z)\\
&-\alpha_D(y)\dashv(x\vdash z)+\alpha_D(y)\dashv(z\dashv x).
\end{aligned}$$

The left hand side vanishes thanks also to the Definition 4.3. Thus the conclusion holds. □

5 Modules over color Hom-Leibniz algebras

In this section we introduce modules over color Hom-Leibniz algebras and prove that the Yau's twisting of module structure map works very well with module over color Hom-Leibniz algebras.

Definition 5.1. *Let G be an abelian group. A Hom-module is a pair (M,α_M) in which M is a G-graded vector space and $\alpha_M : M\longrightarrow M$ is an even linear map.*

Definition 5.2. *Let $(L,[-,-],\varepsilon,\alpha_L)$ be a color Hom-Leibniz algebra and (M,α_M) a Hom-module. An L-module on M consists of even $\mathbf{K}$-bilinear maps $\mu_L : L\otimes M\to M$ and $\mu_R : M\otimes L\to M$ such that for any $x,y\in L$, and $m\in M$,*

$$\begin{aligned}
\alpha_M(\mu_L(x,m)) &= \mu_L(\alpha_L(x),\alpha_M(m)) && (14)\\
\alpha_M(\mu_R(m,x)) &= \mu_R(\alpha_M(m),\alpha_L(x)) && (15)\\
\mu_L([-,-]\otimes\alpha_M) &= \mu_L(\alpha_L\otimes\mu_L)-\varepsilon(x,y)\mu_L(\alpha_L\otimes\mu_L)(\tau_{L,L}\otimes Id_L), && (16)\\
\mu_R(\alpha_M\otimes[-,-]) &= \mu_R(\mu_L\otimes\alpha_L)(\tau_{M,L}\otimes Id_L)+\varepsilon(x,m)\mu_L(\alpha_L\otimes\mu_R)(\tau_{M,L}\otimes Id_L), && (17)\\
\mu_L(\alpha_L\otimes\mu_R) &= \mu_R(\mu_L\otimes\alpha_L)+\varepsilon(m,x)\mu_R(\alpha_M\otimes[-,-])(\tau_{L,M}\otimes Id_L). && (18)
\end{aligned}$$

where $\tau_{L,L}(x\otimes y)=y\otimes x$, $\tau_{L,M}(x\otimes m)=m\otimes x$, $\tau_{M,L}(m\otimes x)=x\otimes m$.

Remark 5.1. *The conditions (16), (17) and (18) can be written respectively as*

$$\begin{aligned}
[x,y]\cdot\alpha_M(m) &= \alpha_L(x)\cdot(y\cdot m)-\varepsilon(x,y)\alpha_L(y)\cdot(x\cdot m), && (19)\\
\alpha_M(m)*[x,y] &= (x\cdot m)*\alpha_L(y)+\varepsilon(x,m)\alpha_L(x)\cdot(m*y), && (20)\\
\alpha_L(x)\cdot(m*y) &= (x\cdot m)*\alpha_L(y)+\varepsilon(m,x)\alpha_M(m)*[x,y], && (21)
\end{aligned}$$

where $\mu(x\otimes y)=xy$, $\mu_L(x\otimes m)=x\cdot m$ and $\mu_R(m\otimes x)=m*x$.

Example 5.1. *Any color Hom-Leibniz algebra and any Hom-Leibniz superalgebras is a module over itself.*

Theorem 5.1. *Let $(L,[-,-],\varepsilon,\alpha_L)$ be a color Hom-Leibniz algebra and (M,μ_L,μ_R,α_M) a color Hom-Leibniz module. Then,*

$$\tilde{\mu}_L = \mu_L(\alpha_L^2\otimes Id_M) : L\otimes M\to M, \tag{22}$$
$$\tilde{\mu}_R = \mu_R(Id_M\otimes\alpha_L^2) : L\otimes M\to M, \tag{23}$$

define another color Hom-Leibniz module structure on M.

Proof. We have to point out relations (16)-(18) for $\tilde{\mu}_L$ and $\tilde{\mu}_R$. We have, respectively, any for $x,y\in L, m\in M$,

$$\begin{aligned}
\tilde{\mu}_L([-,-]\otimes\alpha_M)(x\otimes y\otimes m) &= \tilde{\mu}_L([x,y]\otimes\alpha_M(m))\\
&= \alpha_L^2([x,y])\cdot\alpha_M(m)\\
&= [\alpha_L^2(x),\alpha_L^2(y)]\cdot\alpha_M(m)\\
&= \alpha_L^3(x)\cdot(\alpha_L^2(y)\cdot m)-\varepsilon(x,y)\alpha_L^3(y)\cdot(\alpha_L^2(x)\cdot m)\ (\text{by (19)})\\
&= \tilde{\mu}_L(\alpha_L\otimes\tilde{\mu}_L)(x\otimes y\otimes m)\\
&\quad -\varepsilon(x,y)\tilde{\mu}_L(\alpha_L\otimes\tilde{\mu}_L)(\tau_{L,L}\otimes Id_L)(x\otimes y\otimes m).
\end{aligned}$$

$$\begin{aligned}
\tilde{\mu}_R(\alpha_M\otimes[-,-])(m\otimes x\otimes y) &= \tilde{\mu}_R(\alpha_M(m)\otimes[x,y])\\
&= \alpha_M(m)*[\alpha_L^2(x),\alpha_L^2(y)]\\
&= (\alpha_L^2(x)\cdot m)*\alpha_L^3(y)+\varepsilon(x,m)\alpha_L^3(x)\cdot(m*\alpha_L^2(y)) \quad (\text{by (20)})\\
&= \tilde{\mu}_R(\tilde{\mu}_L\otimes\alpha_L)(\tau_{M,L}\otimes Id_L)(m\otimes x\otimes y)\\
&\quad +\varepsilon(x,m)\tilde{\mu}_L(\alpha_L\otimes\tilde{\mu}_R)(\tau_{M,L}\otimes Id_L)(m\otimes x\otimes y).
\end{aligned}$$

$$\begin{aligned}
\varepsilon(m,x)\tilde{\mu}_R(\alpha_M\otimes[-,-])(\tau_{L,M}\otimes Id_L)(x\otimes m\otimes y) &= \varepsilon(m,x)\tilde{\mu}_R(\alpha_M\otimes[-,-])(x\otimes m\otimes y)\\
&= \varepsilon(m,x)\tilde{\mu}_R(\alpha_M(m)\otimes[x,y])\\
&= \varepsilon(m,x)\alpha_M(m)*[\alpha_L^2(x),\alpha_L^2(y)]\\
&= \alpha_L^3(x)\cdot(m*\alpha_L^2(y))\\
&\quad -(\alpha_L^2(x)\cdot m)*\alpha_L^3(y) \quad (\text{by (21)})\\
&= \tilde{\mu}_L(\alpha_L\otimes\tilde{\mu}_R)(x\otimes m\otimes y)\\
&\quad -\tilde{\mu}_R(\tilde{\mu}_L\otimes\alpha_L)(x\otimes m\otimes y).
\end{aligned}$$

□

Corollary 5.1. *Let $A_{\alpha_A^n}=(A,[-,-]_{\alpha_A^n},\varepsilon,\alpha_A^n)$ be a multiplicative color Hom-Leibniz algebra as in Theorem 4.1 and (M,μ_L,μ_R,α_M) an $A_{\alpha_A^n}$-module. Then $(M,\tilde{\mu}_L,\tilde{\mu}_R,\alpha_M)$ is also a module over $A_{\alpha_A^n}$.*

References

[1] Abdaoui K., Mabroukand S., Makhlouf A., *Cohomology of Hom-Leibniz and n-ary Hom-Nambu-Lie superalgebras,* 1406.3776v1, 14 Jun 2014 .

[2] Abdaoui K., Ammar F. and Makhlouf A., *Constructions and cohomology of color Hom-Lie algebras*, arXiv :13072612v1, 9 jul 2013.

[3] Aizawa N. and Sato H., *q-deformation of the Virasoro algebra with central extension*, Physics Letters B 256 (1991), 185-190.

[4] Akivis M. A., *Local algebras of multidimensionsal three-web*, Siberian Math. J. **17** (1976), no 1, 3-8.

[5] Akivis M. A., *Three web of multidimensionsal surfaces*, Trudy Geo. Sem. **2** (1969), 7-31 (Russian).

[6] Ammar F., Ayadi I., Mabrouk S. and Makhlouf A., *Quadratique color Hom-Lie algebras*, arXiv 1204.5155, 23 April 2012 .

[7] Ammar F., Mabrouk S., and Makhlouf A., *Quadratic n-ary Hom-Nambu algebras*, (2011), e-Print: arXiv:1110.2078.

[8] Ayupov Sh. A. and Omirov B.A., *On Leibniz algebras*, in Algebras and Operators Theory, Procceding of the Colloquim in Tashkent , Kluver (1998) 1-13.

[9] Bakayoko I., *Modules over color Hom-Poisson algebras*, to appear in Journ. of Gen. Lie Theory and Appl.

[10] Bakayoko, I., *L-modules, L-comodules and Hom-Lie quasi-bialgebras*, African Diaspora Journal of Mathematics, Vol 17 (2014), 49-64.

[11] Bremner M., *Algebras, dialgebras and polynomial identities*, Serdica Math. J. **38** (2012), 91-136.

[12] Cacas J. M., M. A. Insua and N. P. Rego, *On universal central extensions of Hom-Leibniz algebras*, arXiv 1209.6266v1, 27 Sept 2012.

[13] Casas J. M., *Trialgebras and Leibniz 3-algebras* Bol. Soc. Mat. Mexicana (3) Vol 12, 2006.

[14] Cheng Y. S. and Su Y. C., *(Co)homology and universal central extensions of Hom-Leibniz algebras*, Acta Math. Sin. (Engl. Ser.) 27 (5) (2011), 813-830.

[15] Elhamdadi M. and Makhlouf A., *Deformations of Hom-Alternative and Hom-Malcev algebras*, (2010), To appear in Group, Algebra and Geometries, e-Print: arXiv:1006.2499.

[16] Hofmann K. H., Strambach K., *Lie's fondamental theorems of local analytical loops*, Pacific J. Math. **123** (1986), no. 2, 301-327.

[17] Hu N., Pei Y. and Liu D., *A cohomological characterizations of Leibniz central extensions of Lie algebras*, Procceding of American Math. Soc., Vol 136, No 2, 437-447, Feb 2008.

[18] Issa A. N., *Some characterizations of Hom-Leibniz algebras*, Int. Elect. Journ. of Alg., 14 (2013), 1-9.

[19] Issa A. N., *Hom-Akivis algebras*, Comment. Math. Uni. Carolin. 52.4, 485-500 (2011).

[20] Loday J.-L and Pirashvili T., *Universal enveloping algebras of Leibniz algebras*, Georgian Math. J. **5** (1998), 263-676.

[21] Loday J.-L., *Une version non-commutative des algbres de Lie : Les algbres de Leibniz*, Enseig. Math., 39 (1993), 269-293.

[22] Makhlouf A., *Hom-alternative algebras and Hom-Jordan algebras*, Int. Electron. J. Algebra 8 (2010), 177-190.

[23] Makhlouf A. and Silvestrov S., *Notes on Formal Deformations of Hom-associative algebras and Hom-Lie algebras*, Forum Math., 22 (2010), no. 4, 715-739.

[24] Makhlouf A. and Silvestrov S., *Hom-algebra structures*, J. Gen. Lie Theory Appl. Vol.2 (2008), no. 2, 51-64.

[25] Miràs J. M. C., *Central extensions of Leibniz algebras*, Extracta Mathematica, Vol **13**, No 3, 393-397 (1998).

[26] Ratseev S. M., *On varities of Leibniz Poisson algebras with the identity* $\{x,y\}\cdot\{z,t\}=0$, Journal of Siberian Federal University . Math. and Phys. 6(1), 97-104 (2013).

[27] Ratseev S. M., *Commutative Leibniz-Poisson algebras of polynomial growth*, Vestn. Samar. Gos. Univ. Estestvennonauch. Ser., **94**(2012), no 3/1, 54-65 (in Russian).

[28] Yau D., *Hom-Malcev, Hom-alternative and Hom-Jordan algebras*, Int. Elect. Journ. of Alg., 11 (2012), 177-217.

[29] Yau D., *Module Hom-algebras*, arXiv:0812.4695v1, 26 Dec 2008.

[30] Yau D., *Envelopping algebras of Hom-Lie algebras* , ArXiv : 0709.0849v1, 6 Sept. 2007.

[31] Yuan L. , *Hom-Lie color algebras*, communication in algebra, **V**ol 40, no. 2, (2012), 575-592.